

10/599,002

Connecting via Winsock to STN

Welcome to STN International! Enter x:x



\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 11:21:18 ON 30 JUL 2009

=> file reg

=>

Uploading C:\Program Files\Stnexp\Queries\11599002.str



chain nodes :  
11 12 13 14 15  
ring nodes :

10/599,002

1 2 3 4 5 6 7 8 9 10  
chain bonds :  
6-14 10-11 11-12 12-13 14-15  
ring bonds :  
1-2 1-6 2-3 2-7 3-4 3-10 4-5 5-6 7-8 8-9 9-10  
exact/norm bonds :  
10-11 11-12 12-13 14-15  
exact bonds :  
6-14  
normalized bonds :  
1-2 1-6 2-3 2-7 3-4 3-10 4-5 5-6 7-8 8-9 9-10  
isolated ring systems :  
containing 1 :

G1:N,Hy

Match level :


1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom  
11:CLASS 12:CLASS 13:CLASS 14:CLASS 15:Atom

L1 STRUCTURE UPLOADED

=> d l1

L1 HAS NO ANSWERS

L1 STR



G1 N,Hy

Structure attributes must be viewed using STN Express query preparation.

=> s l1 sam

SAMPLE SEARCH INITIATED 11:22:19 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 68 TO ITERATE

100.0% PROCESSED 68 ITERATIONS

2 ANSWERS

SEARCH TIME: 00.00.01


10/599,002

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*  
BATCH \*\*COMPLETE\*\*  
PROJECTED ITERATIONS: 866 TO 1854  
PROJECTED ANSWERS: 2 TO 124

L2 2 SEA SSS SAM L1

=> d scan

L2 2 ANSWERS REGISTRY COPYRIGHT 2009 ACS on STN  
IN Quinoline, 3-(phenylsulfonyl)-8-[3-(1-piperidinyl)propoxy]-, hydrochloride  
(1:1)  
MF C23 H26 N2 O3 S . Cl H



● HCl

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):end

=> s l1 full  
FULL SEARCH INITIATED 11:22:27 FILE 'REGISTRY'  
FULL SCREEN SEARCH COMPLETED - 1471 TO ITERATE

100.0% PROCESSED 1471 ITERATIONS 30 ANSWERS  
SEARCH TIME: 00.00.01

L3 30 SEA SSS FUL L1


=> file ca

=> s l3  
L4 1 L3

=> d ibib abs fhitstr

L4 ANSWER 1 OF 1 CA COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 143:386931 CA  
 TITLE: Preparation of  
 3-[(hetero)arylsulfonyl]-8-[(aminoalkyl)oxy]quinolines  
 as 5-HT6 receptor antagonists for the treatment of CNS  
 disorders  
 INVENTOR(S): Ahmed, Mahmood; Johnson, Christopher Norbert; Miller,  
 Neil Derek; Trani, Giancarlo; Witty, David R.  
 PATENT ASSIGNEE(S): Glaxo Group Limited, UK; Witty, David R  
 SOURCE: PCT Int. Appl., 34 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005095346	A1	20051013	WO 2005-GB1106	20050324
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1730112	A1	20061213	EP 2005-729157	20050324
EP 1730112	B1	20080903		
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, HR, LV				
JP 2007530648	T	20071101	JP 2007-505618	20050324
AT 407120	T	20080915	AT 2005-729157	20050324
ES 2313319	T3	20090301	ES 2005-729157	20050324
US 20070191345	A1	20070816	US 2006-599002	20060918
PRIORITY APPLN. INFO.:			GB 2004-7025	A 20040329
			WO 2005-GB1106	W 20050324
OTHER SOURCE(S):			CASREACT 143:386931; MARPAT 143:386931	
GI				



AB The title compds. I [R1 = (un)substituted NH2 or a N containing heterocyclyl;  
X = a bond, (un)substituted CH2, (CH2)2, etc.; R2 = halo, CN, CF3, etc.; n  
= 0-3; R3, R4 = H, halo, CN, etc.; A = (un)substituted (hetero)aryl,  
arylaryl, etc.], useful in the treatment of CNS and other disorders, were  
prepared Thus, reacting 2-dimethylaminoethanol with  
3-phenylsulfonyl-8-iodoquinoline (preparation given) afforded 48% I [R1 = NMe2;  
X = (CH2)2; R2-R4 = H; A = Ph] which was converted to its HCl salt which  
showed antagonist potency for the 5-HT6 receptor, having fpKi > 8.0 at  
human cloned 5-HT6 receptors. The pharmaceutical composition comprising the  
compound I is disclosed.

IT 866782-65-2P  
RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic  
preparation); THU (Therapeutic use); BIOL (Biological study); PREP  
(Preparation); RACT (Reactant or reagent); USES (Uses)  
(preparation of 3-[(hetero)arylsulfonyl]-8-[(aminoalkyl)oxy]quinolines as  
5-HT6 receptor antagonists for the treatment of CNS disorders)

RN 866782-65-2 CA  
CN Quinoline, 3-(phenylsulfonyl)-8-[2-(1-pyrrolidinyloxy)]- (CA INDEX  
NAME)



OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD  
(3 CITINGS)

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> file marpat  
COST IN U.S. DOLLARS  
FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
5.85	191.95

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)  
CA SUBSCRIBER PRICE

SINCE FILE	TOTAL
ENTRY	SESSION
-0.78	-0.78

FILE 'MARPAT' ENTERED AT 11:22:40 ON 30 JUL 2009  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

10/599,002

COPYRIGHT (C) 2009 American Chemical Society (ACS)

FILE CONTENT: 1961-PRESENT VOL 151 ISS 4 (20090724/ED)

MARPAT RECORDS ARE DERIVED FROM INPI DATA FOR 1961-1987

MOST RECENT CITATIONS FOR PATENTS FROM MAJOR ISSUING AGENCIES  
(COVERAGE TO THESE DATES IS NOT COMPLETE):

US 20090149676 11 JUN 2009  
DE 102007059214 10 JUN 2009  
EP 2065708 03 JUN 2009  
JP 2009137851 25 JUN 2009  
WO 2009074534 18 JUN 2009  
GB 2453808 22 APR 2009  
FR 2924713 12 JUN 2009  
RU 2357978 10 JUN 2009  
CA 2643394 07 MAY 2009

The new MARPAT User Guide is now available at:  
<http://www.cas.org/support/stngen/stdoc/marpat.html>.

=> s l3 full  
FULL SEARCH INITIATED 11:22:45 FILE 'MARPAT'  
FULL SCREEN SEARCH COMPLETED - 8854 TO ITERATE

100.0% PROCESSED 8854 ITERATIONS 14 ANSWERS  
SEARCH TIME: 00.00.04

L5 14 SEA SSS FUL L1

=> d ibib abs fqhit 1-14

L5 ANSWER 1 OF 14 MARPAT COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 149:425659 MARPAT  
TITLE: Preparation of phenyl-naphthalenols and related  
compounds as (17) $\beta$ -hydroxy steroid dehydrogenase  
inhibitors  
INVENTOR(S): Hartmann, Rolf; Frotscher, Martin; Oberwinkler,  
Sandrine; Ziegler, Erika; Messinger, Josef; Thole,  
Heinrich-Hubert  
PATENT ASSIGNEE(S): Universitaet des Saarlandes, Germany  
SOURCE: PCT Int. Appl., 125pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: German  
FAMILY ACC. NUM. COUNT: 2  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2008116920	A2	20081002	WO 2008-EP53672	20080327
WO 2008116920	A3	20090402		
W: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,				

ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH,  
 PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM,  
 TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW  
 RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU,  
 IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK,  
 TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,  
 TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW,  
 AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA


DE 102007015169 A1 20081002 DE 2007-10200701516920070327

PRIORITY APPLN. INFO.: DE 2007-10200701516920070327

GI



I



II

AB Title compds. I [W, X, Y, Z = C=, N=; R1 = H, halo, OH, etc.; R2 = H, halo, OH, etc.; R3 = H, halo, OH, etc.; R4 = H, OH; R5 = H, halo, OH, etc.; R6 = H, halo, OH, etc.; R7 = H, halo, OH, etc.; R8 = H, halo, OH, etc. R9 = H, OH, CN, etc.; R10 = H, OH, CN, etc.] and their pharmaceutically acceptable salts and formulations were prepared For example, Suzuki coupling of 6-bromo-2-naththol and phenylboronic acid afforded claimed phenylnaphthalenol II in 87% yield. In (17) $\beta$ -hydroxysteroid dehydrogenase 1 inhibition assays, 17-examples of compds. I exhibited IC50 values ranging from 7-840 nM.

MSTR 1

G4—G19

G4 = 45


G12—G10  
45

G5 = bond


G10 = heteroaryl <containing up to 12 atoms,  
 1 or more heteroatoms, zero or more N, zero or more O,  
 zero or more S (no other heteroatoms)> (opt. substd.) / Ph

G12 = SO2


G19 = 7



G22 = 108 / 115



G32 = 268 / N



Patent location: claim 1  
 Note: and pharmacologically acceptable salts  
 Note: substitution is restricted  
 Note: also incorporates claim 11

L5 ANSWER 2 OF 14 MARPAT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 148:419118 MARPAT

TITLE: Measurement of rate of light-induced change of dye optical property in presence of probe-target hybrids for detecting polynucleotides

INVENTOR(S): Bupp, Charles Robert; Choi, K. Yeon; Koshinsky, Heather; Nulf, Christopher; Urdea, Mickey; Wang, Miaomiao; Zwick, Michael

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 66pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20080096193	A1	20080424	US 2006-362359	20061024
PRIORITY APPLN. INFO.:			US 2006-362359	20061024

AB Methods of determining the presence or amount of a target polynucleotide in a sample are provided. A sample that contains a target polynucleotide, a nucleic acid analog that is complementary to a target nucleic acid sequence of the target polynucleotide, and a cyanine dye for which the rate of change in an optical property is different in the presence and absence of a target polynucleotide/nucleic acid analog hybrid are combined to produce a reaction mixture. The rate of change in an optical property of the dye in the reaction mixture is compared to a reference value characteristic of the rate of change in the optical property of the dye in a similar reaction mixture containing a known amount of a polynucleotide/nucleic acid analog.



hybrid to determine a relative rate of change in the optical property. The relative rate of change in the optical property of dye in the reaction mixture is correlated with the presence or amount of the specified target polynucleotide in the sample.

MSTR 1




G1 = 31 / 33 / 38

$\begin{matrix} \text{G10-G6} \\ 31 \end{matrix}$       $\begin{matrix} \text{G7-G10-G19} \\ 33 \end{matrix}$       $\begin{matrix} \text{G9-G22} \\ 38 \end{matrix}$

G3 = 58 / 60 / 65

$\begin{matrix} \text{G10-G17} \\ 58 \end{matrix}$       $\begin{matrix} \text{G18-G10-G20} \\ 60 \end{matrix}$       $\begin{matrix} \text{G9-G22} \\ 65 \end{matrix}$

G4 = bond  
G5 = 54-17 55-19



G9 = SO2  
G10 = O  
G11 = heteroaryl <containing 1 or more heteroatoms, zero or more N, zero or more O, zero or more S (no other heteroatoms), mono- or bicyclic>  
G17 = carbon chain <containing 1-9 C, 0 or more double bonds, 0 or more triple bonds> (opt. substd. by 1 or more G11)  
G22 = Ph (opt. substd.)  
Patent location: claim 1  
Note: or salts or esters  
Note: additional interruption also claimed  
Note: substitution is restricted

L5 ANSWER 3 OF 14 MARPAT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 147:358263 MARPAT

TITLE: Carbocyanine dye dimers linked by a conjugated alkenyl chain for use in detection of nucleic acid

INVENTOR(S): hybridization  
 Bupp, Charles R., II; Choi, K. Yeon; Holmes-Davis,  
 Rachel Anne; Izmailov, Alexander; Koshinsky, Heather;  
 Nulf, Christopher J.; Urdea, Micky; Wang, Miaomiao;  
 Warner, Brian David; Zwick, Michael

PATENT ASSIGNEE(S): Investigen, Inc., USA

SOURCE: PCT Int. Appl., 229 pp.  
 CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English


FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2007100711	A2	20070907	WO 2007-US4814	20070223
WO 2007100711	A3	20090409		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA			
US 20070231821	A1	20071004	US 2007-710667	20070223
EP 2010677	A2	20090107	EP 2007-751566	20070223
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR			
PRIORITY APPLN. INFO.:			US 2006-776595P	20060224
			WO 2007-US4814	20070223

AB Dimeric carbocyanine dyes linked by a conjugated alkenyl moiety that change optical properties upon binding nucleic acids are described for use in quant. hybridization assays. The dyes can be used as reporters in assays using nucleic acid probes, or with analogs such as peptide nucleic acids or locked nucleic acids as probes. The rate of change in an optical property of the dye in the hybridization is compared to a reference value characteristic of the rate of change in the optical property of the dye in a similar reaction mixture containing a known quantity of a to determine a relative rate of change in the optical property. The relative rate of change in the optical property of dye in the reaction mixture is correlated with the presence or amount of the specified target polynucleotide in the sample. The dyes are hydrophobic and a detergent is necessary for their solubilization. Optimization of assay conditions and the determination of sensitivities of assays using different conditions, probe types, light sources and analyte sequence concns. are reported.

MSTR 1




G1 = 31 / 33 / 38

<sup>G10-G6</sup><sub>31</sub>      <sup>G7-G10-G19</sup><sub>33</sub>      <sup>G9-G22</sup><sub>38</sub>

G3 = 58 / 60 / 65

<sup>G10-G17</sup><sub>58</sub>      <sup>G18-G10-G20</sup><sub>60</sub>      <sup>G9-G22</sup><sub>65</sub>

G4 = bond  
G5 = 54-17 55-19




G9 = SO2  
G10 = O  
G11 = heteroaryl <containing 1 or more heteroatoms,  
zero or more N, zero or more O,  
zero or more S (no other heteroatoms), mono- or bicyclic>  
G17 = carbon chain <containing 1-9 C,  
0 or more double bonds, 0 or more triple bonds>  
(opt. substd. by 1 or more G11)  
G22 = Ph (opt. substd.)  
Patent location: claim 1  
Note: or salts or esters  
Note: additional interruption also claimed  
Note: substitution is restricted

L5 ANSWER 4 OF 14 MARPAT COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 146:358716 MARPAT  
TITLE: Preparation of alkyl sulfoxide quinolines as Nk-3  
receptor ligands  
INVENTOR(S): Albert, Jeffrey S.; Koether, Gerard M.; Alhambra,  
Cristobal; Kang, James; Simpson, Thomas R.; Woods,  
James; Li, Yan  
PATENT ASSIGNEE(S): Astrazeneca AB, Swed.  
SOURCE: PCT Int. Appl., 68pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent

LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2007035158	A1	20070329	WO 2006-SE1068	20060919
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
AU 2006292849	A1	20070329	AU 2006-292849	20060919
CA 2621062	A1	20070329	CA 2006-2621062	20060919
EP 1928835	A1	20080611	EP 2006-784189	20060919
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, RS			
JP 2009508946	T	20090305	JP 2008-532190	20060919
MX 2008003765	A	20080402	MX 2008-3765	20080318
KR 2008046669	A	20080527	KR 2008-706742	20080320
IN 2008DN02393	A	20080725	IN 2008-DN2393	20080320
NO 2008001862	A	20080616	NO 2008-1862	20080417
US 20080214605	A1	20080904	US 2008-67572	20080421
CN 101312950	A	20081126	CN 2006-80043132	20080519
PRIORITY APPLN. INFO.:			US 2005-719275P	20050921
			US 2005-719286P	20050921
			WO 2006-SE1068	20060919


GI



AB Title compds. represented by the formula I [wherein R1 = H, (cyclo)alkyl or alkyl-OCO-; A = Ph or cycloalkyl; R2, R3 = independently H, OH, NH2, etc.; n = 1-3; R4 = E-SOr-(CH2)p-; E = (cyclo)alkyl or (hetero)aryl; p = 0-6; r = 0-2; R5 = independently H, halo, amino, etc.; q = 1-3; and

stereoisomer, enantiomer, in vivo-hydrolyzable precursor or pharmaceutically acceptable salts thereof] were prepared as NK-3 receptor ligands. For example, amidation of 3-(methylthio)-2-phenylquinoline-4-carboxylic acid with ((S)-(-)-1-phenylpropyl)amine gave II in 70% yield. The biol. test for NK-3 receptor binding activity was described (no data). I and their pharmaceutical compns. are useful for the treatment or prophylaxis of a disease or condition in which modulation of the NK-3 receptor is beneficial, such as mental and behavioral disorders (no data).

MSTR 1



G4 = 51 / 299

$$\begin{matrix} \text{G12-G13-G14} & \text{G13-G15} \\ 51 & 299 \end{matrix}$$

G5 = 75

$$\begin{matrix} \text{G18-G9} \\ 75 \end{matrix}$$

G9 = alkyl <containing 1-6 C>  
(opt. substd. by 1 or more G10)

G10 = NH2

G12 = (0-6) CH2

G13 = SO2

G14 = 4-pyridyl

G18 = O

Patent location:

claim 1

Note:

or in vivo hydrolysable precursors or  
pharmaceutically acceptable salts

Stereochemistry:

or stereoisomers or enantiomers

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 5 OF 14 MARPAT COPYRIGHT 2009 ACS on STN


ACCESSION NUMBER: 145:377221 MARPAT

TITLE: Preparation of dihydro pyridines, quinolines, and

isoquinolines as anti-Alzheimer agents.  
 INVENTOR(S): Marsais, Francis; Bohn, Pierre; Levacher, Vincent; Le Fur, Nicolas  
 PATENT ASSIGNEE(S): Insa Rouen, Fr.; Gous Inc.  
 SOURCE: PCT Int. Appl., 136pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006103120	A2	20061005	WO 2006-EP3787	20060329
WO 2006103120	A3	20070215		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM EP 1731507 A1 20061213 EP 2005-290914 20050426 R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, LV, MK, YU AU 2006228683 A1 20061005 AU 2006-228683 20060329 CA 2603345 A1 20061005 CA 2006-2603345 20060329 EP 1868998 A2 20071226 EP 2006-742673 20060329 R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR IN 2007DN07518 A 20071109 IN 2007-DN7518 20070928 US 20090062279 A1 20090305 US 2008-909911 20080716 PRIORITY APPLN. INFO.: EP 2005-290719 20050401 EP 2005-290914 20050426 WO 2006-EP3787 20060329				


OTHER SOURCE(S): CASREACT 145:377221  
 GI




AB Title compds. [I; dotted lines = double bond between CR5-CR6, and another double bond between either CR2CR3 or CR3CR4; R1-R6 = H, OH, alkyl, aryl,

heteroaryl, aralkyl, alkylaryl, alkoxy, hydroxyalkyl, alkoxyalkyl, Ph, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>H, Z, Z<sub>1</sub>; R<sub>4</sub>R<sub>5</sub>, R<sub>5</sub>R<sub>6</sub> = atoms to form (substituted) 6-membered aryl, 5-6 membered heterocyclyl; ≥1 of R<sub>2</sub>, R<sub>3</sub>, R<sub>5</sub> = CO<sub>2</sub>R, COSR, CONRR', cyano, COR, CF<sub>3</sub>, SOR, SO<sub>2</sub>R, SONRR', SO<sub>2</sub>NRR', NO<sub>2</sub>, halo, heteroaryl; R, R' = H, alkyl, cycloalkyl, aminoalkyl, aryl,, heteroaryl, etc.; NRR' = (substituted) heterocyclyl; Z = LmZ<sub>1</sub>; L = alkyl, aryl, heteroaryl, Ph, alkylaryl, aralkyl; Z<sub>1</sub> = XC(:Y)NR<sub>7</sub>R<sub>8</sub>; X, Y = O, S; R<sub>7</sub>, R<sub>8</sub> = H, alkyl, aryl, heteroaryl, aralkyl, Ph, cyclopropyl, (CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>H; n = 1-6; with a proviso], were prepared Thus, Et 1-methyl-7-N,N-dimethylcarbamoyloxy-1,4-dihydroquinoline-3-carboxylate (5-step preparation from 3-cyano-7-methoxyquinoline given) inhibited human acetylcholinesterase with IC<sub>50</sub> = 0.5 μM.


MSTR 1




G1 = 5



G2 = 289-2 290-3 288-4 282-6 287-7




G3 = N  
G4 = 39



G11 = SO<sub>2</sub>  
G12 = 169

10/599,002



G13 = O  
G14 = NH2  
G17 = 166

G11-G12  
166

G34 = 339

G11-G12  
339

Patent location: claim 1  
Note: G9 is optionally present  
Note: substitution is restricted  
Note: or pharmaceutically acceptable salts  
Note: also incorporates claim 2, structure G+ and claim 40, structures E1, E2, and E3  
Stereochemistry: or stereoisomers

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 6 OF 14 MARPAT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 144:88181 MARPAT

TITLE: Heteroaryl sulfones and sulfonamides and the preparation, pharmaceutical compositions, and therapeutic uses thereof, particularly for treatment of proliferative diseases such as cancer

INVENTOR(S): Reddy, Premkumar E.; Reddy, Ramana M. V.

PATENT ASSIGNEE(S): Temple University-of the Commonwealth System of Higher Education, USA

SOURCE: PCT Int. Appl., 136 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005123077	A2	20051229	WO 2005-US20023	20050608
WO 2005123077	A3	20060526		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU,



ZA, ZM, ZW  
 RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,  
 AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,  
 EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT,  
 RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML,  
 MR, NE, SN, TD, TG  
 AU 2005253966 A1 20051229 AU 2005-253966 20050608  
 CA 2569705 A1 20051229 CA 2005-2569705 20050608  
 EP 1765804 A2 20070328 EP 2005-763374 20050608  
 R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,  
 IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA,  
 HR, LV, MK, YU  
 JP 2008501804 T 20080124 JP 2007-527659 20050608  
 IN 2006DN07044 A 20070713 IN 2006-DN7044 20061123  
 US 20070232649 A1 20071004 US 2006-628019 20061128  
 MX 2006014230 A 20070214 MX 2006-14230 20061206  
 KR 2007034574 A 20070328 KR 2007-700456 20070108  
 PRIORITY APPLN. INFO.: US 2004-578162P 20040608  
 WO 2005-US20023 20050608  
 GI


\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB Title compds. I [wherein: R1 = halo, hydrocarbyl, acyl, various  
 disubstituted amino or substituted hydroxy derivs., NO<sub>2</sub>, cyano, PO<sub>3</sub>H<sub>2</sub>  
 diesters, mono- or disubstituted SO<sub>2</sub>NH<sub>2</sub>, guanidino or monohydrocarbyl  
 derivs., haloalkyl, or heteroalkyl; R2 = (un)substituted (hetero)aryl; M =  
 bond, NR<sub>3</sub>, CH:CH, CHR<sub>4</sub>, CH(R<sub>4</sub>)A(CH:CH)<sub>m</sub>; N(R<sub>3</sub>)A(CH:CH)<sub>m</sub>; R<sub>3</sub>, R<sub>4</sub> = H,  
 alkyl; A = SO<sub>2</sub>, CO; Q = O, S, or NH; n = 0-4; m = 0-1; with provisos;  
 including salts] are disclosed. I are useful as antiproliferative agents  
 including, for example, as anticancer agents. Examples include approx. 60  
 prepared compds. I and 2 bioassays. Over 350 invention compds. are also  
 named in claims. For instance, cyclocondensation of  
 5-bromosalicylaldehyde with [(4-methoxyanilino)sulfonyl]acetic acid in  
 refluxing AcOH in the presence of PhCH<sub>2</sub>NH<sub>2</sub> gave invention compound II. In  
 tests against 5 human tumor cell lines, invention compound III had 50%  
 growth-inhibitory concns. (IG<sub>50</sub>) of 12-16 μM.

MSTR 1


G4—SO<sub>2</sub>—G24  
 109

G1 = O / NH  
 G2 = 199




10/599,002

G4 = 8



G24 = 123



G32 = alkylene <containing 2-6 C>

Patent location: claim 1

Note: additional ring formation also claimed

Note: substitution is restricted

Note: or salts

Note: also incorporates claim 47 and 48

L5 ANSWER 7 OF 14 MARPAT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 143:386931 MARPAT

TITLE: Preparation of  
3-[(hetero)arylsulfonyl]-8-[(aminoalkyl)oxy]quinolines  
as 5-HT<sub>6</sub> receptor antagonists for the treatment of CNS  
disorders

INVENTOR(S): Ahmed, Mahmood; Johnson, Christopher Norbert; Miller,  
Neil Derek; Trani, Giancarlo; Witty, David R.

PATENT ASSIGNEE(S): Glaxo Group Limited, UK; Witty, David R

SOURCE: PCT Int. Appl., 34 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005095346	A1	20051013	WO 2005-GB1106	20050324
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,				


EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT,  
 RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML,  
 MR, NE, SN, TD, TG

EP 1730112	A1	20061213	EP 2005-729157	20050324
EP 1730112	B1	20080903		

R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,  
 IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, HR, LV


JP 2007530648	T	20071101	JP 2007-505618	20050324
AT 407120	T	20080915	AT 2005-729157	20050324
ES 2313319	T3	20090301	ES 2005-729157	20050324
US 20070191345	A1	20070816	US 2006-599002	20060918
PRIORITY APPLN. INFO.:			GB 2004-7025	20040329
			WO 2005-GB1106	20050324

OTHER SOURCE(S): CASREACT 143:386931  
 GI



AB The title compds. I [R1 = (un)substituted NH2 or a N containing heterocyclyl;  
 X = a bond, (un)substituted CH2, (CH2)2, etc.; R2 = halo, CN, CF3, etc.; n  
 = 0-3; R3, R4 = H, halo, CN, etc.; A = (un)substituted (hetero)aryl,  
 arylaryl, etc.], useful in the treatment of CNS and other disorders, were  
 prepared. Thus, reacting 2-dimethylaminoethanol with  
 3-phenylsulfonyl-8-iodoquinoline (preparation given) afforded 48% I [R1 = NMe2;  
 X = (CH2)2; R2-R4 = H; A = Ph] which was converted to its HCl salt which  
 showed antagonist potency for the 5-HT6 receptor, having fpKi > 8.0 at  
 human cloned 5-HT6 receptors. The pharmaceutical composition comprising the  
 compound I is disclosed.


MSTR 1




G1 = G5  
 G2 = NH2

10/599,002

G5 = (1-3) 14



G8 = Ph  
G25 = 200



Patent location: claim 1

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L5 ANSWER 8 OF 14 MARPAT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 139:265380 MARPAT  
TITLE: Hair dye compositions containing quinolinium salts  
INVENTOR(S): Sauter, Guido; Braun, Hans-Juergen; Duc-Reichlin, Nadia  
PATENT ASSIGNEE(S): Wella Aktiengesellschaft, Germany  
SOURCE: Eur. Pat. Appl., 14 pp.  
CODEN: EPXXDW  
DOCUMENT TYPE: Patent  
LANGUAGE: German  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1346719	A1	20030924	EP 2002-25423	20021115
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
DE 10211413	A1	20030925	DE 2002-10211413	20020315
US 20030177592	A1	20030925	US 2003-361380	20030210
US 6977001	B2	20051220		
BR 2003000496	A	20040810	BR 2003-496	20030313
PRIORITY APPLN. INFO.:			DE 2002-10211413	20020315

AB The invention concerns hair dyes that are prepared from two components; component A1 contains a quinolinium derivative; component A2 includes a nucleophile compound Other direct dyes can be added; solns., emulsions, creams, foams, gels can be formulated. Thus component A1 contained (g): 4-chloro-1-ethylquinolinium tetrafluoroborate 0.70 decyl glycoside 4.0; EDTA disodium salt 0.2; ethanol 5.0; water to 100. Component A2 included: 1,4-diaminobenzene 0.27; decyl glycoside 4.0; EDTA disodium salt 0.2; ethanol 5.0; 25% ammonia solution 6.0; water to 100.

MSTR 1



G2 = 27 / 52



G3 = tolyl

G5 = heteroaryl (opt. substd.)

G7 = O

G14 = SO2

Patent location: claim 1

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 9 OF 14 MARPAT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 138:271682 MARPAT

TITLE: Preparation of cyclic hydroxamic acids as inhibitors of matrix metalloproteinases and/or TNF- $\alpha$  converting enzyme for treatment of inflammatory disorders

INVENTOR(S): Ott, Gregory; Chen, Xiao-Tao; Duan, Jingwu; Lu, Zhonghui

PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA

SOURCE: PCT Int. Appl., 344 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003024899	A2	20030327	WO 2002-US29685	20020916
WO 2003024899	A3	20031127		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				



AU 2002341715	A1	20030401	AU 2002-341715	20020916
US 20030139388	A1	20030724	US 2002-244626	20020916
US 6740649	B2	20040525		
EP 1427408	A2	20040616	EP 2002-775865	20020916

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK

PRIORITY APPLN. INFO.:

US 2001-322630P	20010917
WO 2002-US29685	20020916

GI




AB Title compds. I [wherein ring B = (un)substituted 4-7 membered (hetero)cyclic ring containing 0-2 O, N, NR1, or SOp atoms and 0-3 carbonyl groups; R1 and R2 = independently Q, alk(en/yn)ylene-Q, or (un)substituted alkylene-Q interrupted by O, NRa, CO, CO2, CONRa, NRaCO, NRaCO2, NRaCONRa, SOp, NRaSO2, or SO2NRa; or R1 = (un)substituted alkylene-Q interrupted by OCO, OCO2, or OCONRa; Q = H or (un)substituted (hetero)cyclyl; R3 = Q1, Cl, F, alk(en/yn)ylene-Q1, or (un)substituted alkylene-Q1 interrupted by O, NR1, NRaCO, CONRa, CO, CO2, SOp, or SO2NRa; Q1 = H or (un)substituted Ph, naphthyl, or heterocyclyl; Za = (un)substituted benzimidazolyl, indolyl, imidazopyridinyl, pyrazolylpyridinyl, benzofuranyl, benzothiazinyl, quinolinyl, etc.; Ra = independently H, alkyl, Ph, or benzyl; p = 0-2; or stereoisomers or pharmaceutically acceptable salts thereof] were prepared as inhibitors of matrix metalloproteinases (MMP), TNF- $\alpha$  converting enzyme (TACE), aggrecanase, or a combination thereof. For example, reaction of benzyl Me maleate with paraformaldehyde and glycine gave benzyl Me (cis)-3,4-pyrrolidinedicarboxylate (100%). BOC-protection (64%), debenzylation (96%), resolution of the (3S,4S)-isomer with (S)- $\alpha$ -methylbenzylamine, conversion to the carbamate with DPPA and PhCH2OH (76%), and Pd catalyzed hydrogenation (100%) provided Me (3S,4S)-4-amino-1-(tert-butoxycarbonyl)-3-pyrrolidinecarboxylate. Coupling of the amine with 4-[(2-methylthio-1H-benzimidazol-1-yl)methyl]benzoic acid (preparation given) afforded the amide (99%), which was treated with NH2OH•HCl/MeONa to give the hydroxamic acid (3S,4S)-II


10/599,002

(33%). A number of the compds. of the invention inhibited MMP-1, 2, 3, 7, 8, 9, 10, 12, 13, 14, 15, and/or 16 with  $K_i$  values of  $\leq 10 \mu\text{M}$ . Thus, I are useful for the treatment of a wide variety of inflammatory disorders (no data).


MSTR 1



G3 = 16




G5 = 80-13 78-44 81-17 82-20



G11 = 111

~~G15-G14~~  
111

G14 = Ph  
G15 = SO<sub>2</sub>  
G17 = O  
G19 = C(O)  
G29 = 212 / 227



Patent location: claim 1  
Note: or pharmaceutically acceptable salts  
Note: substitution is restricted  
Note: additional ring formation also claimed  
Stereochemistry: or stereoisomers

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 10 OF 14 MARPAT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 138:78134 MARPAT  
 TITLE: Direct hair dyes composed of 1-benzopyrane-derivatives  
 and an electrophilic substance  
 INVENTOR(S): Sauter, Guido; Braun, Hans-Juergen; Brouillard,  
 Raymond; Fougrousse, Andre; Roehri-Stoeckel,  
 Christine  
 PATENT ASSIGNEE(S): Wella Aktiengesellschaft, Germany  
 SOURCE: PCT Int. Appl., 51 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
WO 2003000214	A1	20030103	WO 2002-EP1194	20020206
W:			AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW	
RW:			GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG	
DE 10130144	A1	20030102	DE 2001-10130144	20010622
AU 2002246084	A1	20030108	AU 2002-246084	20020206
BR 2002005662	A	20030715	BR 2002-5662	20020206
EP 1404289	A1	20040407	EP 2002-714147	20020206
R:			AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR	
JP 2004521144	T	20040715	JP 2003-506861	20020206
US 20030196281	A1	20031023	US 2003-380896	20030320
PRIORITY APPLN. INFO.:			DE 2001-10130144	20010622
			WO 2002-EP1194	20020206


AB The invention concerns a two component hair dye where the components are mixed in the presence of acids or bases if required to form a direct dye that can be removed with sulfite-containing reducing agents if required. The first component includes 1-benzopyrane-derivs.; the second component contains an electrophilic substance that is selected from the group of carbonyls, imines and 1-alkyl-quinoline derivs. Thus a first components was composed of (g): 7-hydroxy-4-methyl-2-phenyl-1-benzylpyrylium chloride 3.14; cetylstearyl alc. 12.0; Brij 78 P 2.8; ethanol 24.8; water to 100. The second component was a mixture of (g): 4-hydroxy-3-methoxy-benzaldehyde 1.75; cetylstearyl alc. 12.0; Brij 78 P 2.8; ethanol 24.8; water to 100.

MSTR 2


G1—G3 G10

G1 = 10





G3 = 47 / 68



G4 = heteroaryl (opt. substd.)

G5 = tolyl

Patent location: claim 1

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 11 OF 14 MARPAT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 134:17502 MARPAT

TITLE: Preparation of phenoxypropylamine compounds as antagonists of 5-HT1A receptor

INVENTOR(S): Nishiyama, Akira; Bougauchi, Masahiro; Kuroita, Takanobu; Minoguchi, Masanori; Morio, Yasunori; Kanzaki, Kouji

PATENT ASSIGNEE(S): Welfide Corp., Japan

SOURCE: PCT Int. Appl., 335 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: Japanese


FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000071517	A1	20001130	WO 2000-JP3279	20000522
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2375008	A1	20001130	CA 2000-2375008	20000522
BR 2000011542	A	20020305	BR 2000-11542	20000522
EP 1188747	A1	20020320	EP 2000-927844	20000522
EP 1188747	B1	20050907		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
HU 2002001540	A2	20020828	HU 2002-1540	20000522

HU 2002001540	A3	20021228		
NZ 516111	A	20030530	NZ 2000-516111	20000522
CN 1164574	C	20040901	CN 2000-808604	20000522
AU 777594	B2	20041021	AU 2000-46160	20000522
AT 303987	T	20050915	AT 2000-927844	20000522
ES 2244438	T3	20051216	ES 2000-927844	20000522
IL 146564	A	20061231	IL 2000-146564	20000522
JP 3893878	B2	20070314	JP 2000-619774	20000522
US 20020111358	A1	20020815	US 2001-990389	20011123
US 6720320	B2	20040413		
MX 2001012046	A	20030904	MX 2001-12046	20011123
KR 799134	B1	20080129	KR 2001-715024	20011123
ZA 2001010137	A	20030225	ZA 2001-10137	20011210
US 20040138227	A1	20040715	US 2003-740418	20031222
US 7196199	B2	20070327		
KR 2007118193	A	20071213	KR 2007-726625	20071115
KR 882544	B1	20090212		
PRIORITY APPLN. INFO.:			JP 1999-142750	19990524
			JP 1999-166160	19990614
			JP 1999-277384	19990929
			JP 2000-18080	20000125
			WO 2000-JP3279	20000522
			KR 2001-715024	20011123
			US 2001-990389	20011123


GI




AB Phenoxypropylamine compds. represented by general formula [I; a bond represented by a solid and a dotted line is a double or single bond; X = H, HO, C1-8 alkoxy, acyloxy, oxo; R1 = 4-substituted piperidino, piperazino, 1-piperidinylamino, or 1,2,3,6-tetrahydropyrazinyl, (un)substituted aryloxy- or arylthioamino, (un)substituted heterocyclyloxy- or heterocyclylthioamino, etc.; R3 = H, C1-18 alkyl, halo; Ra, Rb, Rc = H, C1-18 alkyl, OH, C1-8 alkoxy, halo, acyl, NO2, NH2], optically active isomers thereof or pharmaceutically acceptable salts thereof and hydrates of the same are prepared. These compds. have an affinity selectively for 5-HT1A receptor, simultaneously show an antagonistic activity, and inhibit the reuptake of 5-HT, thereby being usable as antidepressant agents quickly achieving an antidepressant effect (no data). Thus, 4-(3,4-dichlorophenyl)piperazine was added to a solution of (S)-5-(4-glycidyoxybenzo[b]furan-2-yl)-3-methylisoxazole in MeOH and refluxed for 8 h to give (S)-1-(4-(3,4-dichlorophenyl)piperazin-1-yl)-3-(2-(3-methylisoxazol-5-yl)benzo[b]furan-4-yloxy)-2-propanol.

MSTR 1A


10/599,002




G3 = 26




G4 = 3




G12 = 176-3 183-1




G14 = 73



G16 = C(O)  
G18 = 75



G19 = SO<sub>2</sub>  
G20 = 83



Patent location: claim 1

Note: and pharmacologically acceptable salts or hydrates

REFERENCE COUNT: 151 THERE ARE 151 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 12 OF 14 MARPAT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 133:89542 MARPAT

TITLE: Preparation of quinoxalines as non-peptide GLP-1 agonists

INVENTOR(S): Teng, Min; Truesdale, Larry Kenneth; Bhumralkar, Dilip; Kiel, Dan; Johnson, Michael D.; Thomas, Christine; Jorgensen, Anker Steen; Madsen, Peter; Olesen, Preben Houlberg; Knudsen, Liselotte Bjerre; Petterson, Ingrid Vivika; Cornelis De Jong, Johannes; Behrens, Carsten; Kodra, Janos Tibor; Lau, Jesper

PATENT ASSIGNEE(S): Novo Nordisk A/S, Den.; Agouron Pharmaceuticals, Inc.

SOURCE: PCT Int. Appl., 194 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent


LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000042026	A1	20000720	WO 2000-DK14	20000114
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
EP 1147094	A1	20011024	EP 2000-900499	20000114
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
JP 2002534512	T	20021015	JP 2000-593594	20000114
US 6927214	B1	20050809	US 2000-483504	20000114
PRIORITY APPLN. INFO.:			DK 1999-41	19990115
			US 1999-116116P	19990115
			WO 2000-DK14	20000114


GI



I

AB The title compds. I [R1, R2, R3, R4 independently = H, halogen, CN, CF3, NO2, OR5, lower alkyl, SR5, S(O2)NR5R6, etc (a proviso is given); A, B = H, halogen, OH, CF3, CF2CF3, CN, NO2, alkyl, alkenyl, etc; L, M = (CH2)mS(CH2)n, (CH2)mO(CH2)n, (CH2)mS(O)(CH2)n, (CH2)mS(O)2(CH2)n, etc; X, V = :N or :CD; D = H, halogen, CN, CF3, NO2, etc; m, n independently = 0, 1, 2, 3, or 4 ] useful as non-peptide GLP-1 agonists for the treatment and/or prevention of disorders and diseases wherein an activation of the human GLP-1 receptor is beneficial, especially metabolic disorders such as Type 1 diabetes, Type 2 diabetes and obesity (no data), are prepared Formulations are given.

MSTR 1



G1 = 15 / 17 / 29 / 34

$\begin{matrix} \text{G3} \\ 15 \end{matrix} \text{---} \text{G2}$      $\begin{matrix} \text{G4} \\ 17 \end{matrix} \text{---} \text{G5}$      $\begin{matrix} \text{G11} \\ 29 \end{matrix} \text{---} \text{G2}$      $\begin{matrix} \text{G6} \\ 34 \end{matrix} \text{---} \text{G2}$

G2 = alkyl <containing 1-6 C>  
(substd. by heterocycle <containing 3-10 atoms,  
1 or more heteroatoms, zero or more N, zero or more O,  
zero or more S (no other heteroatoms), non-aromatic,  
0 or more double-exact bonds> (opt. substd.))

G3 = 0  
G5 = 19

$\begin{matrix} \text{G6} \\ 19 \end{matrix} \text{---} \text{G2}$


G7 = 1 or more N / 23

$\begin{matrix} \text{C} \\ 23 \end{matrix} \text{---} \text{G8}$

G13 = 42

$\begin{matrix} \text{G14} \\ 42 \end{matrix} \text{---} \text{G15}$

G14 = SO2  
G15 = 203



Patent location: claim 1

REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 13 OF 14 MARPAT COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 120:191707 MARPAT

TITLE: 2-Substituted saccharin derivative proteolytic enzyme inhibitors

INVENTOR(S): Hlasta, Dennis John; Desai, Ranjit Chimanlal; Subramanyam, Chakrapani; Lodge, Eric Piatt; Dunlap, Richard Paul; Boaz, Neil Warren; Mura, Albert Joseph; Latimer, Lee Hamilton

PATENT ASSIGNEE(S): Sterling Winthrop Inc., USA

SOURCE: Eur. Pat. Appl., 77 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent


LANGUAGE: English

FAMILY ACC. NUM. COUNT: 7

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 542372	A1	19930519	EP 1992-203469	19921112
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
US 5236917	A	19930817	US 1991-793033	19911115
AU 9225340	A	19930520	AU 1992-25340	19920925
AU 654581	B2	19941110		
CA 2079822	A1	19930516	CA 1992-2079822	19921005
NO 9204401	A	19930518	NO 1992-4401	19921113
NO 303119	B1	19980602		
HU 66873	A2	19950130	HU 1992-3566	19921113
IL 103748	A	19970218	IL 1992-103748	19921113
RU 2101281	C1	19980110	RU 1992-4381	19921113
JP 05194444	A	19930803	JP 1992-305295	19921116
US 5371074	A	19941206	US 1993-67637	19930524
US 5650422	A	19970722	US 1994-270964	19940705
US 5596012	A	19970121	US 1995-449152	19950524
US 5874432	A	19990223	US 1997-803297	19970220
PRIORITY APPLN. INFO.:			US 1991-793033	19911115
			US 1989-347125	19890504
			US 1989-347126	19890504
			US 1990-514920	19900426
			US 1993-67637	19930524
			US 1994-270964	19940705

GI




AB The title compds. I [L = O, S, SO, SO<sub>2</sub>; R<sub>1</sub> = (un)substituted Ph, (un)substituted heterocyclyl, etc.; R<sub>2</sub> = H, lower alkoxy, carbonyl, Ph, PhS; R<sub>3</sub> = H, halogen, (un)substituted alkyl, Ph, lower alkoxy, lower alkoxy, carbonyl, CN, etc.; R<sub>4</sub> = H or 1-3 substituents selected from halogen, CN, NO<sub>2</sub>, NH<sub>2</sub>, etc.; m, n = 0, 1; when m = 0 then R<sub>1</sub> can only be heterocyclyl and CHR<sub>2</sub> can only be bonded to a ring N of R<sub>1</sub>; when m = 0, n = 1 and L is O, S, or SO, then R<sub>2</sub>-R<sub>4</sub> = H; when m = 0, n = 1, L is S, R<sub>2</sub>, R<sub>4</sub> = H and R<sub>3</sub> = halogen; when m = 0, n = 1, and L is SO or SO<sub>2</sub> then R<sub>2</sub> is lower alkoxy, carbonyl and R<sub>3</sub> = R<sub>4</sub> = H while R<sub>1</sub> ≠ substituted Ph], useful for the treatment of degenerative diseases (no data), are prepared. Thus, 2-hydroxymethyl-4-chlorosaccharin was reacted with thionyl chloride, producing 2-chloromethyl-4-chlorosaccharin (II). II demonstrated inhibition constant for human leukocyte elastase (rate of reactivation of enzyme to rate of inactivation of enzyme) of 0.5 nM and 26 nM for α-chymotrypsin.

MSTR 1A



G1 = O  
 G2 = bond  
 G3 = 163



G12 = 81

O<sub>2</sub>S—G13—G14  
 81

G13 = phenylene

Patent location: claim 1


10/599,002

Note: substitution is restricted

L5 ANSWER 14 OF 14 MARPAT COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 119:213908 MARPAT  
TITLE: Silver halide photographic material  
INVENTOR(S): Fukuwa, Junichi; Kobayashi, Akira; Goto, Kenji  
PATENT ASSIGNEE(S): Konica Co., Japan  
SOURCE: Can. Pat. Appl., 71 pp.  
CODEN: CPXXEB  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
CA 2065106	A1	19921005	CA 1992-2065106	19920403
JP 05197057	A	19930806	JP 1992-110787	19920403
PRIORITY APPLN. INFO.:			JP 1991-99626	19910404

GI



AB A Ag halide photog. material for high-contrast dot image formation is disclosed. The material comprises a support and provided thereon a Ag halide emulsion layer and layers adjacent to the emulsion layer. The emulsion is subjected to desalinization comprising using denatured gelatin in the process of preparation thereof. At least one of the layers contains a hydrazine derivative and a compound selected from the group consisting of those represented by formulas A(CH<sub>2</sub>)<sub>n</sub>SC(:N+HR<sub>1</sub>)NHR<sub>1</sub> X<sup>-</sup> (A = OH, SO<sub>3</sub><sup>-</sup>, or N(R<sub>2</sub>)<sub>2</sub>; R<sub>1</sub> = H, (substituted) alkyl having 1-5 C atoms, or (substituted) Ph; R<sub>2</sub> = (substituted) alkyl having 1-5 C atoms; X<sup>-</sup> = an anion), (R<sub>3</sub>)<sub>2</sub>N(CH<sub>2</sub>)<sub>n</sub>SC(S)N(R<sub>4</sub>)<sub>2</sub> (R<sub>3</sub> = H, (substituted) alkyl having 1-5 C atoms, or (substituted) aryl; R<sub>4</sub> = (substituted) alkyl having 1-5 C atoms or (substituted) Ph; n = an integer of 2-5), or I (Q = a group of atoms necessary to form a 5- or 6-membered heterocyclic ring which may be condensed with a benzene or heterocyclic ring; M = H, an alkali metal atom, an ammonium group, or an amine residue).


MSTR 3B

G1—G2


G1 = 232



10/599,002



G7 = SO<sub>2</sub>Ph / 43





Patent location: claim 1

=> file reg

=>

Uploading C:\Program Files\Stnexp\Queries\12599002.str

10/599,002



```
chain nodes :  
11 12 13 16 17  
ring nodes :  
1 2 3 4 5 6 7 8 9 10  
chain bonds :  
6-12 10-11 11-16 12-13 16-17  
ring bonds :  
1-2 1-6 2-3 2-7 3-4 3-10 4-5 5-6 7-8 8-9 9-10  
exact/norm bonds :  
10-11 11-16 12-13 16-17  
exact bonds :  
6-12  
normalized bonds :  
1-2 1-6 2-3 2-7 3-4 3-10 4-5 5-6 7-8 8-9 9-10  
isolated ring systems :  
containing 1 :
```

10/599,002

G1:N,Hy

Match level :


1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom  
11:CLASS 12:CLASS 13:Atom 16:Atom 17:CLASS

L6 STRUCTURE UPLOADED

=> d 16

L6 HAS NO ANSWERS

L6 STR



G1 N,Hy

Structure attributes must be viewed using STN Express query preparation.

=> s 16 sam

SAMPLE SEARCH INITIATED 11:24:40 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 289 TO ITERATE

100.0% PROCESSED 289 ITERATIONS

1 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*

BATCH \*\*COMPLETE\*\*

PROJECTED ITERATIONS: 4761 TO 6799

PROJECTED ANSWERS: 1 TO 80


L7 1 SEA SSS SAM L6

=> d scan

10/599,002

L7 1 ANSWERS REGISTRY COPYRIGHT 2009 ACS on STN  
IN 3-Furanamine, tetrahydro-4-[[3-(phenylsulfonyl)-8-quinolinyl]oxy]-,  
hydrochloride (1:1), (3S,4R)-  
MF C19 H18 N2 O4 S . Cl H

Absolute stereochemistry.



● HCl

ALL ANSWERS HAVE BEEN SCANNED

=> s 16 full  
FULL SEARCH INITIATED 11:24:49 FILE 'REGISTRY'  
FULL SCREEN SEARCH COMPLETED - 5502 TO ITERATE

100.0% PROCESSED 5502 ITERATIONS 4 ANSWERS  
SEARCH TIME: 00.00.02

L8 4 SEA SSS FUL L6

=> file ca

=> s 18  
L9 1 L8

=> d ibib

L9 ANSWER 1 OF 1 CA COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 143:386931 CA  
TITLE: Preparation of  
3-[(hetero)arylsulfonyl]-8-[(aminoalkyl)oxy]quinolines  
as 5-HT<sub>6</sub> receptor antagonists for the treatment of CNS  
disorders  
INVENTOR(S): Ahmed, Mahmood; Johnson, Christopher Norbert; Miller,  
Neil Derek; Trani, Giancarlo; Witty, David R.  
PATENT ASSIGNEE(S): Glaxo Group Limited, UK; Witty, David R  
SOURCE: PCT Int. Appl., 34 pp.

CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005095346	A1	20051013	WO 2005-GB1106	20050324
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1730112	A1	20061213	EP 2005-729157	20050324
EP 1730112	B1	20080903		
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, HR, LV				
JP 2007530648	T	20071101	JP 2007-505618	20050324
AT 407120	T	20080915	AT 2005-729157	20050324
ES 2313319	T3	20090301	ES 2005-729157	20050324
US 20070191345	A1	20070816	US 2006-599002	20060918
PRIORITY APPLN. INFO.:			GB 2004-7025	A 20040329
			WO 2005-GB1106	W 20050324
OTHER SOURCE(S):		CASREACT 143:386931; MARPAT 143:386931		
OS.CITING REF COUNT:	3	THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)		
REFERENCE COUNT:	7	THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT		

=> d his

(FILE 'HOME' ENTERED AT 11:21:18 ON 30 JUL 2009)

FILE 'REGISTRY' ENTERED AT 11:22:04 ON 30 JUL 2009

L1 STRUCTURE UPLOADED  
 L2 2 S L1 SAM  
 L3 30 S L1 FULL

FILE 'CA' ENTERED AT 11:22:29 ON 30 JUL 2009

L4 1 S L3

FILE 'MARPAT' ENTERED AT 11:22:40 ON 30 JUL 2009

L5 14 S L3 FULL

FILE 'REGISTRY' ENTERED AT 11:24:23 ON 30 JUL 2009

L6 STRUCTURE UPLOADED  
 L7 1 S L6 SAM  
 L8 4 S L6 FULL



10/599,002

FILE 'CA' ENTERED AT 11:24:53 ON 30 JUL 2009  
L9 1 S L8

=> file reg

=>

Uploading C:\Program Files\Stnexp\Queries\13599002.str



chain nodes :  
11 12 13 14  
ring nodes :  
1 2 3 4 5 6 7 8 9 10  
chain bonds :  
6-13 10-11 11-12 13-14  
ring bonds :  
1-2 1-6 2-3 2-7 3-4 3-10 4-5 5-6 7-8 8-9 9-10  
exact/norm bonds :  
10-11 11-12 13-14  
exact bonds :  
6-13  
normalized bonds :

10/599,002

1-2 1-6 2-3 2-7 3-4 3-10 4-5 5-6 7-8 8-9 9-10  
isolated ring systems :  
containing 1 :

G1:N,Hy

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom  
11:CLASS 12:CLASS 13:CLASS 14:Atom

L10 STRUCTURE UPLOADED

=> d 1109


L109 NOT FOUND

The L-number entered has not been defined in this session, or it  
has been deleted. To see the L-numbers currently defined in this  
session, enter DISPLAY HISTORY at an arrow prompt (=>).

=> d 110

L10 HAS NO ANSWERS

L10 STR



G1 N,Hy

Structure attributes must be viewed using STN Express query preparation.

=> s 110 full

FULL SEARCH INITIATED 11:26:04 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 5502 TO ITERATE

100.0% PROCESSED 5502 ITERATIONS

26 ANSWERS

SEARCH TIME: 00.00.01

L11 26 SEA SSS FUL L10

=> file ca

=> s 111

10/599,002

L12 1 L11

=> d ibib

L12 ANSWER 1 OF 1 CA COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 143:386931 CA  
TITLE: Preparation of  
3-[(hetero)arylsulfonyl]-8-[(aminoalkyl)oxy]quinolines  
as 5-HT6 receptor antagonists for the treatment of CNS  
disorders  
INVENTOR(S): Ahmed, Mahmood; Johnson, Christopher Norbert; Miller,  
Neil Derek; Trani, Giancarlo; Witty, David R.  
PATENT ASSIGNEE(S): Glaxo Group Limited, UK; Witty, David R  
SOURCE: PCT Int. Appl., 34 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005095346	A1	20051013	WO 2005-GB1106	20050324
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1730112	A1	20061213	EP 2005-729157	20050324
EP 1730112	B1	20080903		
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, HR, LV			
JP 2007530648	T	20071101	JP 2007-505618	20050324
AT 407120	T	20080915	AT 2005-729157	20050324
ES 2313319	T3	20090301	ES 2005-729157	20050324
US 20070191345	A1	20070816	US 2006-599002	20060918
PRIORITY APPLN. INFO.:			GB 2004-7025	A 20040329
			WO 2005-GB1106	W 20050324
OTHER SOURCE(S):	CASREACT 143:386931; MARPAT 143:386931			
OS.CITING REF COUNT:	3	THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)		
REFERENCE COUNT:	7	THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT		

=> d his

(FILE 'HOME' ENTERED AT 11:21:18 ON 30 JUL 2009)

FILE 'REGISTRY' ENTERED AT 11:22:04 ON 30 JUL 2009

L1 STRUCTURE UPLOADED



10/599,002

L2                2 S L1 SAM  
L3                30 S L1 FULL

FILE 'CA' ENTERED AT 11:22:29 ON 30 JUL 2009  
L4                1 S L3

FILE 'MARPAT' ENTERED AT 11:22:40 ON 30 JUL 2009  
L5                14 S L3 FULL

FILE 'REGISTRY' ENTERED AT 11:24:23 ON 30 JUL 2009  
L6                STRUCTURE UPLOADED  
L7                1 S L6 SAM  
L8                4 S L6 FULL

FILE 'CA' ENTERED AT 11:24:53 ON 30 JUL 2009  
L9                1 S L8

FILE 'REGISTRY' ENTERED AT 11:25:14 ON 30 JUL 2009  
L10               STRUCTURE UPLOADED  
L11               26 S L10 FULL

FILE 'CA' ENTERED AT 11:26:06 ON 30 JUL 2009  
L12               1 S L11

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

INTERNATIONAL LOGOFF AT 11:26:19 ON 30 JUL 2009